



US007452966B2

(12) **United States Patent**
Glaesner et al.

(10) **Patent No.:** **US 7,452,966 B2**
(45) **Date of Patent:** ***Nov. 18, 2008**

(54) **GLP-1 ANALOG FUSION PROTEINS**

(75) Inventors: **Wolfgang Glaesner**, Indianapolis, IN (US); **Rohn Lee Millican, Jr.**, Indianapolis, IN (US); **Andrew Mark Vick**, Fishers, IN (US)

(73) Assignee: **Eli Lilly and Company**, Indianapolis, IN (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 26 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **10/558,627**

(22) PCT Filed: **Jun. 10, 2004**

(86) PCT No.: **PCT/US2004/015595**

§ 371 (c)(1),
(2), (4) Date: **Nov. 29, 2005**

(87) PCT Pub. No.: **WO2005/000892**

PCT Pub. Date: **Jan. 6, 2005**

(65) **Prior Publication Data**

US 2007/0036806 A1 Feb. 15, 2007

Related U.S. Application Data

(60) Provisional application No. 60/477,880, filed on Jun. 12, 2003.

(51) **Int. Cl.**
A61K 38/26 (2006.01)
C07K 14/605 (2006.01)

(52) **U.S. Cl.** **530/324**; 514/12; 530/308

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

FOREIGN PATENT DOCUMENTS

WO WO 02/46227 6/2002

OTHER PUBLICATIONS

STN registry entry, 923950-08-7, Mar. 1, 2007, 2 pages.*
Issacs et al. "A Therapeutic Human IgG4 Monoclonal Antibody That Depletes Target Cells In Humans." Clin. Exp. Immunology, vol. 106, pp. 427-433, 1996.

* cited by examiner

Primary Examiner—Anish Gupta

Assistant Examiner—Ronald T Niebauer

(74) *Attorney, Agent, or Firm*—Gregory A. Cox

(57) **ABSTRACT**

The invention provides specific GLP-1 analogs fused to specific IgG4-Fc derivatives. These fusion proteins have an increased half-life, decreased immunogenicity, and reduce effector activity. The fusion proteins are useful in treating diabetes, obesity, irritable bowel syndrome and other conditions that would be benefited by lowering plasma glucose, inhibiting gastric and/or intestinal motility and inhibiting gastric and/or intestinal emptying, or inhibiting food intake.

5 Claims, No Drawings

US 7,452,966 B2

1

GLP-1 ANALOG FUSION PROTEINS

This is the national phase application, under 35 USC 371, for PCT/US04/015595, filed Jun. 10, 2004, which claims the priority of U.S. provisional application No. 60/477,880, filed Jun. 12, 2003.

FIELD OF THE INVENTION

The present invention relates to glucagon-like peptide analogs fused to proteins that have the effect of extending the in vivo half-life of the peptides. These fusion proteins can be used to treat diabetes as well as a variety of other conditions or disorders.

Glucagon-like peptide-1 (GLP-1) analogs and derivatives show promise in clinical trials for the treatment of type 2 diabetes. GLP-1 induces numerous biological effects such as stimulating insulin secretion, inhibiting glucagon secretion, inhibiting gastric emptying, inhibiting gastric motility or intestinal motility, and inducing weight loss. A significant characteristic of GLP-1 is its ability to stimulate insulin secretion without the associated risk of hypoglycemia that is seen when using insulin therapy or some types of oral therapies that act by increasing insulin expression.

The usefulness of therapy involving GLP-1 peptides has been limited by the fact that GLP-1(1-37) is poorly active, and the two naturally occurring truncated peptides, GLP-1(7-37) OH and GLP-1(7-36)NH₂, are rapidly cleared in vivo and have extremely short in vivo half-lives. It is known that endogenously produced dipeptidyl-peptidase IV (DPP-IV) inactivates circulating GLP-1 peptides by removing the N-terminal histidine and alanine residues and is a major reason for the short in vivo half-life.

Various approaches have been undertaken to extend the elimination half-life of a GLP-1 peptide or reduce clearance of the peptide from the body while maintaining biological activity. One approach involves fusing a GLP-1 peptide to the Fc portion of an immunoglobulin. Immunoglobulins typically have long circulating half-lives in vivo. For example, IgG molecules can have a half-life in humans of up to 23 days. The Fc portion of the immunoglobulin is responsible, in part, for this in vivo stability. GLP-1-Fc fusion proteins take advantage of the stability provided by the Fc portion of an immunoglobulin while preserving the biological activity of the GLP-1 molecule.

Although this approach is feasible for GLP-1 therapeutics (See WO 02/46227), there is a general concern regarding the antigenicity of various fusion proteins when administered repeatedly over prolonged periods of time. This is especially a concern for GLP-1-Fc fusion therapeutics as a patient with diabetes must be treated for her entire life once diagnosed with the disease. In addition, Fc fusion protein therapeutics can be a concern if the Fc portion retains unwanted effector functions.

The present invention seeks to overcome the problems associated with the potential immunogenicity and effector activity associated with administration of GLP-1-Fc fusions by identifying specific GLP-1-Fc fusion proteins that have a reduced risk of inducing an immune response after repeated and prolonged administration and no longer have effector function. These specific fusion proteins have substitutions at various positions in the GLP-1 portion as well as the Fc portion of the molecule. The substitutions described herein provide increased potency, increased in vivo stability, elimination of effector function and decreased likelihood the molecule will be recognized by the adaptive elements of the immune system.

2

Compounds of the present invention include heterologous fusion proteins comprising a GLP-1 analog comprising a sequence selected from the group consisting of

a) His-Xaa₈-Glu-Gly-Thr-Phe-Thr-Ser- (SEQ ID NO:1)

Asp-Val-Ser-Ser-Tyr-Leu-Glu-Glu-

Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-

Trp-Leu-Val-Lys-Gly-Gly-Gly

wherein Xaa₈ is selected from Gly and Val;

b) His-Xaa₈-Glu-Gly-Thr-Phe-Thr-Ser- (SEQ ID NO:2)

Asp-Val-Ser-Ser-Tyr-Leu-Glu-Glu-

Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-

Trp-Leu-Lys-Asn-Gly-Gly-Gly

wherein Xaa₈ is selected from Gly and Val;

c) His-Xaa₈-Glu-Gly-Thr-Phe-Thr-Ser- (SEQ ID NO:3)

Asp-Val-Ser-Ser-Tyr-Leu-Glu-Glu-

Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-

Trp-Leu-Val-Lys-Gly-Gly-Pro

wherein Xaa₈ is selected from Gly and Val;

d) His-Xaa₈-Glu-Gly-Thr-Phe-Thr-Ser- (SEQ ID NO:4)

Asp-Val-Ser-Ser-Tyr-Leu-Glu-Glu-

Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-

Trp-Leu-Lys-Asn-Gly-Gly-Pro

wherein Xaa₈ is selected from Gly and Val;

e) His-Xaa₈-Glu-Gly-Thr-Phe-Thr-Ser- (SEQ ID NO:5)

Asp-Val-Ser-Ser-Tyr-Leu-Glu-Glu-

Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-

Trp-Leu-Val-Lys-Gly-Gly

wherein Xaa₈ is selected from Gly and Val;

f) His-Xaa₈-Glu-Gly-Thr-Phe-Thr-Ser- (SEQ ID NO:6)

Asp-Val-Ser-Ser-Tyr-Leu-Glu-Glu-

Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-

Trp-Leu-Lys-Asn-Gly-Gly

US 7,452,966 B2

3

wherein Xaa₈ is selected from Gly and Val;
fused to the Fc portion of an immunoglobulin comprising
the sequence of SEQ ID NO:7

Ala-Glu-Ser-Lys-Tyr-Gly-Pro-Pro-Cys- (SEQ ID NO:7)
Pro-Pro-Cys-Pro-Ala-Pro-Xaa₁₆-Xaa₁₇-
Xaa₁₈-Gly-Gly-Pro-Ser-Val-Phe-Leu-
Phe-Pro-Pro-Lys-Pro-Lys-Asp-Thr-Leu-
Met-Ile-Ser-Arg-Thr-Pro-Glu-Val-Thr-
Cys-Val-Val-Val-Asp-Val-Ser-Gln-Glu-
Asp-Pro-Glu-Val-Gln-Phe-Asn-Trp-Tyr-
Val-Asp-Gly-Val-Glu-Val-His-Asn-Ala-
Lys-Thr-Lys-Pro-Arg-Glu-Glu-Gln-Phe-
Xaa₈₀-Ser-Thr-Tyr-Arg-Val-Val-Ser-
Val-Leu-Thr-Val-Leu-His-Gln-Asp-Trp-
Leu-Asn-Gly-Lys-Glu-Tyr-Lys-Cys-Lys-
Val-Ser-Asn-Lys-Gly-Leu-Pro-Ser-Ser-
Ile-Glu-Lys-Thr-Ile-Ser-Lys-Ala-Lys-
Gly-Gln-Pro-Arg-Glu-Pro-Gln-Val-Tyr-
Thr-Leu-Pro-Pro-Ser-Gln-Glu-Glu-Met-
Thr-Lys-Asn-Gln-Val-Ser-Leu-Thr-Cys-
Leu-Val-Lys-Gly-Phe-Tyr-Pro-Ser-Asp-
Ile-Ala-Val-Glu-Trp-Glu-Ser-Asn-Gly-
Gln-Pro-Glu-Asn-Asn-Tyr-Lys-Thr-Thr-
Pro-Pro-Val-Leu-Asp-Ser-Asp-Gly-Ser-
Phe-Phe-Leu-Tyr-Ser-Arg-Leu-Thr-Val-
Asp-Lys-Ser-Arg-Trp-Gln-Glu-Gly-Asn-
Val-Phe-Ser-Cys-Ser-Val-Met-His-Glu-
Ala-Leu-His-Asn-His-Tyr-Thr-Gln-Lys-
Ser-Leu-Ser-Leu-Ser-Leu-Gly-Xaa₂₃₀

wherein:

Xaa at position 16 is Pro or Glu;

Xaa at position 17 is Phe, Val, or Ala;

Xaa at position 18 is Leu, Glu, or Ala;

Xaa at position 80 is Asn or Ala; and

Xaa at position 230 is Lys or is absent.

The C-terminus of the GLP-1 analog portion and the N-terminus of the Fc portion of the heterologous fusion proteins of the present invention are preferably fused together via 1, 1.5 or 2 repeats of a G-rich peptide linker having the sequence Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser (SEQ ID NO:8).

The present invention also includes polynucleotides encoding the heterologous fusion proteins of the present invention, as well as vectors and host cells comprising such polynucleotides. Methods of treating patients suffering from non-insulin dependent as well as insulin dependent diabetes mellitus, obesity, and various other disorders and conditions

4

comprising administering the heterologous fusion proteins discussed herein are also encompassed by the present invention.

The heterologous fusion proteins of the present invention comprise a GLP-1 analog portion and an Fc portion. The GLP-1 analog portion and the Fc portion comprise substitutions to the native GLP-1 sequence and the human IgG4 sequence respectively that provide the protein with increased potency and in vivo stability compared to native GLP-1 or GLP-1 analogs not fused to an Fc sequence while decreasing the potential for inducing antibody formation after prolonged and repeated administration in humans.

Native GLP-1 is processed in vivo such that the first 6 amino acids are cleaved from the molecule. Thus, by custom in the art, the amino terminus of GLP-1 has been assigned the number 7 and the carboxy-terminus, number 37. The other amino acids in the polypeptide are numbered consecutively as shown in SEQ ID NO:9. For example, position 8 is alanine and position 22 is glycine. The processed peptide may be further modified in vivo such that the C-terminal glycine residue is removed and replaced with an amide group. Thus, GLP-1(7-37)OH and GLP-1(7-36)amide represent the two native forms of the molecule. GLP-1(7-37)OH has the amino acid sequence of SEQ ID NO:9:

⁷His-Ala-Glu-¹⁰Gly-Thr-Phe-Thr-Ser- (SEQ ID NO:9)
¹⁵Asp-Val-Ser-Ser-Tyr-²⁰Leu-Glu-Gly-
Gln-Ala-²⁵Ala-Lys-Glu-Phe-Ile-³⁰Ala-
Trp-Leu-Val-Lys-³⁵Gly-Arg-³⁷Gly

The GLP-1 analog portion of the heterologous fusion protein comprises three primary substitutions at positions 8, 22, and 36 relative to native GLP-1(7-37). The substitution at position 8 reduces the rate at which the endogenous enzyme dipeptidyl-peptidase IV (DPP-IV) inactivates the analog. DPP-IV cleaves native GLP-1 between the 2nd and 3rd amino acids (between position 8 and 9) and the resulting molecule is less active. Thus, the heterologous fusion proteins of the present invention are DPP-IV resistant. The substitution at position 22 reduces the potential of the molecule to aggregate and increases the potency of the molecule. The substitution at position 36 in the context of the analog with changes at 8 and 22 as well as in the context of the entire fusion protein reduces the risk that the fusion protein will induce a neutralizing immune response after repeated and prolonged administration in humans.

The central event in the generation of both humoral and cell-mediated immune responses is the activation and clonal expansion of T-helper (T_H) cells. T_H cell activation is initiated by interaction of the T-cell receptor (TCR)-CD3 complex with a processed antigenic peptide bound to a class II major histocompatibility (MHC) molecule in the presence of an antigen-presenting cell (APC). Interaction of a T_H cell with antigen initiates a cascade of biochemical events that induces the resting T_H cell to enter the cell cycle (G₀ to G₁ transition). The activated T cell progresses through the cell cycle, proliferating and differentiating into memory cells or effector cells.

US 7,452,966 B2

5

The following sequence was analyzed to identify potential epitopes:

His-Gly-Glu-Gly-Thr-Phe-Thr-Ser-Asp- (SEQ ID NO:10) 5
Val-Ser-Ser-Tyr-Leu-Glu-Glu-Gln-Ala-
Ala-Lys-Glu-Phe-Ile-Ala-Trp-Leu-Val-
Lys-Gly-Arg-Gly-Gly-Gly-Gly-Gly-Ser-
Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-
Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-
Gly-Ser-Gly-Gly-Gly-Gly-Ser-Ala-Glu-
Ser-Lys-Tyr-Gly-Pro-Pro-Cys-Pro

This sequence is a GLP-1 analog sequence with changes at positions 8 and 22 relative to the native sequence followed by 2 copies of a G-rich linker sequence followed by the first 10 amino acids of an Fc region derived from human IgG4. Epitope as used herein refers to a region of a protein molecule to which an antibody can bind. An immunogenic epitope is defined as the part of the protein that elicits an antibody response when the whole protein is the immunogen. Epitope mapping involved the scanning of sequences using a sliding nine amino acid window coupled with advanced statistical analysis techniques to extract the information contained in these patterns. A proprietary software package known as Epi-Matrix™ was used to analyze the sequence and identify peptides that are highly likely to provoke an immune response when presented to T-cells. Eight highly common alleles were used in the analysis for Class II MHC receptor interaction. These alleles included DRB1*0101, DRB1*0301, DRB1*0401, DRB1*0701, DRB1*0801, DRB1*1101, DRB1*1301, and DRB1*1501.

A strong epitope was predicted to be located at the junction of the C-terminus of the GLP-1 analog portion and the beginning of the linker. The sequence of this epitope is Trp-Leu-Val-Lys-Gly-Arg-Gly-Gly-Gly (SEQ ID NO:11) which interacts with DRB1*0801. The present invention encompasses the discovery that this epitope can be eliminated by changing the GLP-1 analog C-terminus to one of the following sequences:

Trp-Leu-Val-Lys-Gly-Gly-Gly; (SEQ ID NO:12)
Trp-Leu-Lys-Asn-Gly-Gly-Gly; (SEQ ID NO:13)
Trp-Leu-Val-Lys-Gly-Gly-Pro; (SEQ ID NO:14) 50
Trp-Leu-Lys-Asn-Gly-Gly-Pro; (SEQ ID NO:15)
Trp-Leu-Val-Lys-Gly-Gly; (SEQ ID NO:16)
and
Trp-Leu-Lys-Asn-Gly-Gly. (SEQ ID NO:17) 55

The heterologous fusion proteins of the present invention contain an Fc portion which is derived from human IgG4, but comprises one or more substitutions compared to the wild-type human sequence. As used herein, the Fc portion of an immunoglobulin has the meaning commonly given to the term in the field of immunology. Specifically, this term refers to an antibody fragment which does not contain the two antigen binding regions (the Fab fragments) from the antibody. The Fc portion consists of the constant region of an antibody from both heavy chains, which associate through non-covalent interactions and disulfide bonds. The Fc portion

6

can include the hinge regions and extend through the CH2 and CH3 domains to the c-terminus of the antibody. The Fc portion can further include one or more glycosylation sites.

There are five types of human immunoglobulins with different effector functions and pharmacokinetic properties. IgG is the most stable of the five types having a serum half-life in humans of about 23 days. There are four IgG subclasses (G1, G2, G3, and G4) each of which have different biological functions known as effector functions. These effector functions are generally mediated through interaction with the Fc receptor (FcγR) or by binding C1q and fixing complement. Binding to FcγR can lead to antibody dependent cell mediated cytotoxicity, whereas binding to complement factors can lead to complement mediated cell lysis. In designing heterologous Fc fusion proteins wherein the Fc portion is being utilized solely for its ability to extend half-life, it is important to minimize any effector function. Thus, the heterologous fusion proteins of the present invention are derived from the human IgG4 Fc region because of its reduced ability to bind FcγR and complement factors compared to other IgG subtypes. IgG4, however, has been shown to deplete target cells in humans [Issacs et al., (1996) Clin. Exp. Immunol. 106:427-433]. Because the heterologous fusion proteins of the present invention target beta cells in the pancreas to induce insulin expression, using an IgG4 derived region in an Fc fusion protein could initiate an immune response against the pancreatic beta cell through interaction of the fusion protein with the GLP-1 receptor present on pancreatic beta cells. Thus, the IgG4 Fc region which is part of the heterologous fusion proteins of the present invention contains substitutions that eliminate effector function. The IgG4 Fc portion of the fusion proteins of the present invention may contain one or more of the following substitutions: substitution of proline for glutamate at residue 233, alanine or valine for phenylalanine at residue 234 and alanine or glutamate for leucine at residue 235 (EU numbering, Kabat, E. A. et al. (1991) *Sequences of Proteins of Immunological Interest*, 5th Ed. U.S. Dept. of Health and Human Services, Bethesda, Md., NIH Publication no. 91-3242). These residues corresponds to positions 16, 17 and 18 in SEQ ID NO:7. Further, removing the N-linked glycosylation site in the IgG4 Fc region by substituting Ala for Asn at residue 297 (EU numbering) which corresponds to position 80 of SEQ ID NO:7 is another way to ensure that residual effector activity is eliminated in the context of a heterologous fusion protein.

In addition, the IgG4 Fc portion of the heterologous fusion proteins of the present invention contain a substitution that stabilizes heavy chain dimer formation and prevents the formation of half-IgG4 Fc chains. The heterologous fusion proteins of the present invention preferably exist as dimers joined together by disulfide bonds and various non-covalent interactions. Wild-type IgG4 contains a Pro-Pro-Cys-Pro-Ser-Cys (SEQ ID NO:18) motif beginning at residue 224 (EU numbering). This motif in a single GLP-1 analog-Fc chain forms disulfide bonds with the corresponding motif in another GLP-1 analog-Fc chain. However, the presence of serine in the motif causes the formation of single chain fusion proteins. The present invention encompasses heterologous Fc fusion proteins wherein the IgG4 sequence is further modified such that serine at position at 228 (EU numbering) is substituted with proline (amino acid residue 11 in SEQ ID NO:7).

The C-terminal lysine residue present in the native molecule may be deleted in the IgG4 derivative Fc portion of the heterologous fusion proteins discussed herein (position 230 of SEQ ID NO:7; deleted lysine referred to as des-K). Fusion proteins expressed in some cell types (such as NS0 cells) wherein lysine is encoded by the C-terminal codon are het-

US 7,452,966 B2

7

erogeneous in that a portion of the molecules have lysine as the C-terminal amino acid and a portion have lysine deleted. The deletion is due to protease action during expression in some types of mammalian cells. Thus, to avoid this heterogeneity, it is preferred that Fc fusion expression constructs lack a C-terminal codon for lysine.

It is preferred that the C-terminal amino acid of the GLP-1 analog portion discussed herein is fused to the N-terminus of the IgG4 Fc analog portion via a glycine-rich linker. The in vivo function and stability of the heterologous fusion proteins of the present invention can be optimized by adding small peptide linkers to prevent potentially unwanted domain interactions. Further, a glycine-rich linker provides some structural flexibility such that the GLP-1 analog portion can interact productively with the GLP-1 receptor on target cells such as the beta cells of the pancreas. These linkers, however, can significantly increase the risk that the fusion protein will be immunogenic in vivo. Thus, it is preferred that the length be no longer than necessary to prevent unwanted domain interactions and/or optimize biological activity and/or stability. The preferred glycine-rich linker comprises the sequence: Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser (SEQ ID NO:8). Although more copies of this linker may be used in the heterologous fusion proteins of the present invention, it is preferred that a single copy of this linker be used to minimize the risk of immunogenicity associated with prolonged and repeated administration.

Preferred GLP-1-Fc heterologous fusion proteins of the present invention include the following proteins: Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-1L-IgG4 (S228P), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-1L-IgG4 (S228P, F234A, L235A), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-1L-IgG4 (S228P, N297A), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-1L-IgG4 (S228P, F234A, L235A, N297A), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-1.5L-IgG4 (S228P), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-1.5L-IgG4 (S228P, F234A, L235A), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-1.5L-IgG4 (S228P, N297A), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-1.5L-IgG4 (S228P, F234A, L235A, N297A), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-2L-IgG4 (S228P), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-2L-IgG4 (S228P, F234A, L235A), Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-2L-IgG4 (S228P, N297A), and Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-2L-IgG4 (S228P, F234A, L235A, N297A), and the Val⁸ and des-K forms of all of the above.

The nomenclature used herein to refer to specific heterologous fusion proteins is defined as follows: Specific substitutions to the GLP-1 portion of the fusion protein are indicated using the specific amino acid being substituted followed by the residue number. GLP-1(7-37) indicates that the GLP-1 portion of the mature fusion protein begins with His at position 7 and ends with Gly at position 37. L refers to a linker with the sequence Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser (SEQ ID NO:8). The number immediately preceding the L refers to the number of linkers separating the GLP-1 portion from the Fc portion. A linker specified as 1.5L refers to the sequence Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser (SEQ ID NO:19). IgG4 refers to an analog of the human IgG4 Fc sequence specified as SEQ ID NO:7. Substitutions in the IgG4 Fc portion of the heterologous fusion protein are indicated in parenthesis. The wild-type amino acid is specified by its common abbreviation followed by the position number in the context of the entire IgG4 sequence using the EU numbering system followed by the amino acid being substituted at that position specified by its common abbreviation.

8

Although the heterologous fusion proteins of the present invention can be made by a variety of different methods, because of the size of the fusion protein, recombinant methods are preferred. For purposes of the present invention, as disclosed and claimed herein, the following general molecular biology terms and abbreviations are defined below.

“Base pair” or “bp” as used herein refers to DNA or RNA. The abbreviations A, C, G, and T correspond to the 5'-monophosphate forms of the deoxyribonucleosides (deoxy)adenosine, (deoxy)cytidine, (deoxy)guanosine, and thymidine, respectively, when they occur in DNA molecules. The abbreviations U, C, G, and A correspond to the 5'-monophosphate forms of the ribonucleosides uridine, cytidine, guanosine, and adenosine, respectively when they occur in RNA molecules. In double stranded DNA, base pair may refer to a partnership of A with T or C with G. In a DNA/RNA, heteroduplex base pair may refer to a partnership of A with U or C with G. (See the definition of “complementary”, *infra*.)

“Digestion” or “Restriction” of DNA refers to the catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA (“sequence-specific endonucleases”). The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors, and other requirements were used as would be known to one of ordinary skill in the art. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer or can be readily found in the literature.

“Ligation” refers to the process of forming phosphodiester bonds between, two double stranded nucleic acid fragments. Unless otherwise provided, ligation may be accomplished using known buffers and conditions with a DNA ligase, such as T4 DNA ligase.

“Plasmid” refers to an extrachromosomal (usually) self-replicating genetic element.

“Recombinant DNA cloning vector” as used herein refers to any autonomously replicating agent, including, but not limited to, plasmids and phages, comprising a DNA molecule to which one or more additional DNA segments can or have been added.

“Recombinant DNA expression vector” as used herein refers to any recombinant DNA cloning vector in which a promoter to control transcription of the inserted DNA has been incorporated.

“Transcription” refers to the process whereby information contained in a nucleotide sequence of DNA is transferred to a complementary RNA sequence.

“Transfection” refers to the uptake of an expression vector by a host cell whether or not any coding sequences are, in fact, expressed. Numerous methods of transfection are known to the ordinarily skilled artisan, for example, calcium phosphate co-precipitation, liposome transfection, and electroporation. Successful transfection is generally recognized when any indication of the operation of this vector occurs within the host cell.

“Transformation” refers to the introduction of DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integration. Methods of transforming bacterial and eukaryotic hosts are well known in the art, many of which methods, such as nuclear injection, protoplast fusion or by calcium treatment using calcium chloride are summarized in J. Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, (1989). Generally, when introducing DNA into Yeast the term transformation is used as opposed to the term transfection.

“Translation” as used herein refers to the process whereby the genetic information of messenger RNA (mRNA) is used to specify and direct the synthesis of a polypeptide chain.

US 7,452,966 B2

9

“Vector” refers to a nucleic acid compound used for the transfection and/or transformation of cells in gene manipulation bearing polynucleotide sequences corresponding to appropriate protein molecules which, when combined with appropriate control sequences, confers specific properties on the host cell to be transfected and/or transformed. Plasmids, viruses, and bacteriophage are suitable vectors. Artificial vectors are constructed by cutting and joining DNA molecules from different sources using restriction enzymes and ligases. The term “vector” as used herein includes Recombinant DNA cloning vectors and Recombinant DNA expression vectors.

“Complementary” or “Complementarity”, as used herein, refers to pairs of bases (purines and pyrimidines) that associate through hydrogen bonding in a double stranded nucleic acid. The following base pairs are complementary: guanine and cytosine; adenine and thymine; and adenine and uracil.

“Primer” refers to a nucleic acid fragment which functions as an initiating substrate for enzymatic or synthetic elongation.

“Promoter” refers to a DNA sequence which directs transcription of DNA to RNA.

“Probe” refers to a nucleic acid compound or a fragment, thereof, which hybridizes with another nucleic acid compound.

“Leader sequence” refers to a sequence of amino acids which can be enzymatically or chemically removed to produce the desired polypeptide of interest.

“Secretion signal sequence” refers to a sequence of amino acids generally present at the N-terminal region of a larger polypeptide functioning to initiate association of that polypeptide with the cell membrane compartments like endoplasmic reticulum and secretion of that polypeptide through the plasma membrane.

Wild-type human IgG4 proteins can be obtained from a variety of sources. For example, these proteins can be obtained from a cDNA library prepared from cells which express the mRNA of interest at a detectable level. Libraries can be screened with probes designed using the published DNA or protein sequence for the particular protein of interest. For example, immunoglobulin light or heavy chain constant regions are described in Adams, et al. (1980) *Biochemistry* 19:2711-2719; Goughet, et al. (1980) *Biochemistry* 19:2702-2710; Dolby, et al. (1980) *Proc. Natl. Acad. Sci. USA* 77:6027-6031; Rice et al. (1982) *Proc. Natl. Acad. Sci. USA* 79:7862-7862; Falkner, et al. (1982) *Nature* 298:286-288; and Morrison, et al. (1984) *Ann. Rev. Immunol.* 2:239-256.

Screening a cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, NY (1989). An alternative means to isolate a gene encoding an immunoglobulin protein is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., *PCR Primer: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, NY (1995)]. PCR primers can be designed based on published sequences.

Generally the full-length wild-type sequences cloned from a particular library can serve as a template to create the IgG4 Fc analog fragments of the present invention that retain the ability to confer a longer plasma half-life on the GLP-1 analog that is part of the fusion protein. The IgG4 Fc analog fragments can be generated using PCR techniques with primers designed to hybridize to sequences corresponding to the desired ends of the fragment. PCR primers can also be designed to create restriction enzyme sites to facilitate cloning into expression vectors.

10

DNA encoding the GLP-1 analogs of the present invention can be made by a variety of different methods including cloning methods like those described above as well as chemically synthesized DNA. Chemical synthesis may be attractive given the short length of the encoded peptide. The amino acid sequence for GLP-1 has been published as well as the sequence of the preproglucagon gene. [Lopez, et al. (1983) *Proc. Natl. Acad. Sci., USA* 80:5485-5489; Bell, et al. (1983) *Nature*, 302:716-718; Heinrich, G., et al. (1984) *Endocrinol*, 115:2176-2181; Ghiglion, M., et al. (1984) *Diabetologia* 27:599-600]. Thus, primers can be designed based on the native sequence to generate DNA encoding the GLP-1 analogs described herein.

The gene encoding a fusion protein can then be constructed by ligating DNA encoding a GLP-1 analog in-frame to DNA encoding the IgG Fc proteins described herein. The DNA encoding wild-type GLP-1 and IgG4 Fc fragments can be mutated either before ligation or in the context of a cDNA encoding an entire fusion protein. A variety of mutagenesis techniques are well known in the art. The gene encoding the GLP-1 analog and the gene encoding the IgG4 Fc analog protein can also be joined in-frame via DNA encoding a G-rich linker peptide. A preferred DNA sequence encoding one of the preferred heterologous fusion proteins of the present invention, Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-1L-IgG4 (S228P, F234A, L235A, des K), is provided as SEQ ID NO:20:

CACGGCGAGGGCACCTTCACCTCCGACGTGTCCTCC (SEQ ID NO:20)

TATCTCGAGGAGCAGGCCGCAAGGAATTCATCGCC

TGGCTGGTGAAGGGCGGCGGCGGTGGTGGTGGCTCC

GGAGGCGGCGGCTCTGGTGGCGGTGGCAGCGCTGAG

TCCAAATATGGTCCCCCATGCCACCCCTGCCAGCA

CCTGAGGCCGCGGGGGACCATCAGTCTTCCTGTTC

CCCCCAAACCCAAGGACACTCTCATGATCTCCCGG

ACCCCTGAGGTACGTCGTCGGTGGTGGACGTGAGC

CAGGAAGACCCCGAGGTCCAGTTCAACTGGTACGTG

GATGGCGTGGAGGTGCATAATGCCAAGACAAAGCCG

CGGGAGGAGCAGTTCAACAGCACGTACCGTGTGGTC

AGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAAC

GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGGC

CTCCCGTCCTCCATCGAGAAAACCATCTCCAAGCC

AAAGGGCAGCCCCGAGAGCCACAGGTGTACACCTG

CCCCCATCCAGGAGGAGATGACCAAGAACCAGGTC

AGCCTGACCTGCCTGGTCAAAGGCTTCTACCCAGC

GACATCGCCGTGGAGTGGGAAAGCAATGGGCAGCCG

GAGAACAACCTACAAGACCACGCCTCCCGTGCTGGAC

TCCGACGGCTCCTTCTTCTCTACAGCAGGCTAACC

GTGGACAAGAGCAGGTGGCAGGAGGGGAATGTCTTC

TCATGCTCCGTGATGCATGAGGCTCTGCACAACCAC

TACACACAGAAGAGCCTCTCCCTGTCTCTGGGT

US 7,452,966 B2

11

Host cells are transfected or transformed with expression or cloning vectors described herein for heterologous fusion protein production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in *Mammalian Cell Biotechnology: A Practical Approach*, M. Butler, ed. (IRL Press, 1991) and Sambrook, et al., supra. Methods of transfection are known to the ordinarily skilled artisan, for example, CaPO₄ and electroporation. General aspects of mammalian cell host system transformations have been described in U.S. Pat. No. 4,399,216. Transformations into yeast are typically carried out according to the method of van Solingen et al., *J Bact.* 130(2): 946-7 (1977) and Hsiao et al., *Proc. Natl. Acad. Sci. USA* 76(8): 3829-33 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene or polyomithine, may also be used. For various techniques for transforming mammalian cells, see Keown, et al., *Methods in Enzymology* 185: 527-37 (1990) and Mansour, et al., *Nature* 336(6197): 348-52 (1988).

Suitable host cells for cloning or expressing the nucleic acid (e.g., DNA) in the vectors herein include yeast or higher eukaryote cells.

Eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for fusion protein vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* [Beach and Nurse, *Nature* 290: 140-3 (1981); EP 139,383 published 2 May 1995]; *Muyveromyces* hosts [U.S. Pat. No. 4,943,529; Fleer, et al., *Bio/Technology* 9(10): 968-75 (1991)] such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574) [de Louvencourt et al., *J. Bacteriol.* 154 (2): 737-42 (1983)]; *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilum* (ATCC 36.906) [Van den Berg et al., *Bio/Technology* 8(2): 135-9 (1990)]; *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070) [Sreekrishna et al., *J. Basic Microbiol.* 28(4): 265-78 (1988)]; *Candida*; *Trichoderma reesia* (EP 244, 234); *Neurospora crassa* [Case, et al., *Proc. Natl. Acad. Sci. USA* 76(10): 5259-63 (1979)]; *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 Oct. 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 Jan. 1991), and *Aspergillus* hosts such as *A. nidulans* [Ballance et al., *Biochem. Biophys. Res. Comm.* 112(1): 284-9 (1983)]; Tilburn, et al., *Gene* 26(2-3): 205-21 (1983); Yelton, et al., *Proc. Natl. Acad. Sci. USA* 81(5): 1470-4 (1984)) and *A. niger* [Kelly and Hynes, *EMBO J.* 4(2): 475-9 (1985)]. Methylo-trophic yeasts are selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotoruia*. A list of specific species that are exemplary of this class of yeast may be found in C. Antony, *The Biochemistry of Methylo-trophs* 269 (1982).

Suitable host cells for the expression of the fusion proteins of the present invention are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sp, *Spodoptera* high5 as well as plant cells. Examples of useful mammalian host cell lines include NSO myeloma cells, Chinese hamster ovary (CHO), SP2, and COS cells. More specific examples include

12

monkey kidney CVI line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line [293 or 293 cells subcloned for growth in suspension culture, Graham, et al., *J. Gen Virol.*, 36(1): 59-74 (1977)]; Chinese hamster ovary cells/-DHFR [CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77(7): 4216-20 (1980)]; mouse sertoli cells [TM4, Mather, *Biol. Reprod.* 23(1):243-52 (1980)]; human lung cells (W138. ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). A preferred cell, line for production of the Fc fusion proteins of the present invention is the NS0 myeloma cell line available from the European Collection of Cell Cultures (ECACC, catalog #85110503) and described in Galfre, G. and Milstein, C. ((1981) *Methods in Enzymology* 73(13):3-46; and *Preparation of Monoclonal Antibodies: Strategies and Procedures*, Academic Press, N.Y., N.Y.).

The fusion proteins of the present invention may be recombinantly produced directly, or as a protein having a signal sequence or other additional sequences which create a specific cleavage site at the N-terminus of the mature fusion protein. In general, the signal sequence may be a component of the vector, or it may be a part of the fusion protein-encoding DNA that is inserted into the vector. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* cc-factor leaders, the latter described in U.S. Pat. No. 5,010, 182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179), or the signal described in WO 90/13646. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., neomycin, methotrexate, or tetracycline, (b) complement autotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the fusion protein-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described [Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77(7): 4216-20 (1980)]. A suitable selection gene for use in yeast is the trp1 gene present in the yeast plasmid YRp7 [Stinchcomb, et al., *Nature* 282(5734): 39-43 (1979); Kingsman, et al., *Gene* 7(2): 141-52 (1979); Tschumper, et al., *Gene* 10(2): 157-66 (1980)]. The trp1 gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEPC1 [Jones, *Genetics* 85: 23-33 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the fusion protein-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Examples of suitable promoting sequences for use with yeast hosts

US 7,452,966 B2

13

include the promoters for 3-phosphoglycerate kinase [Hitze-
man, et al., *J. Biol. Chem.* 255(24): 12073-80 (1980)] or other
glycolytic enzymes [Hess et al., *J. Adv. Enzyme Reg.* 7: 149
(1968); Holland, *Biochemistry* 17(23): 4900-7 (1978)], such
as enolase, glyceraldehyde-3-phosphate dehydrogenase, 5
hexokinase, pyruvate decarboxylase, phosphofructokinase,
glucose-6-phosphate isomerase, 3-phosphoglycerate mutase,
pyruvate kinase, triosephosphate isomerase, phosphoglucose
isomerase, and glucokinase. Other yeast promoters, which
are inducible promoters having the additional advantage of
transcription controlled by growth conditions, are the pro-
moter regions for alcohol dehydrogenase 2, isocytchrome C,
acid phosphatase, degradative enzymes associated with nitro-
gen metabolism, metallothionein, glyceraldehyde-3-phos-
phate dehydrogenase, and enzymes responsible for maltose 10
and galactose utilization. Suitable vectors and promoters for
use in yeast expression are further described in EP 73,657.
Transcription of fusion protein-encoding mRNA from vec-
tors in mammalian host cells may be controlled, for example,
by promoters obtained from the genomes of viruses such as
polyoma virus, fowlpox virus, adenovirus (such as Adenovi-
rus 2), bovine papilloma virus, avian sarcoma virus, cyto-
me-galovirus, a retrovirus, hepatitis-B virus and Simian Virus 40
(SV40), from heterologous mammalian promoters, e.g., the
actin promoter or an immunoglobulin promoter, and from 25
heat-shock promoters, provided such promoters are compat-
ible with the host cell systems.

Transcription of a polynucleotide encoding a fusion pro-
tein by higher eukaryotes may be increased by inserting an
enhancer sequence into the vector. Enhancers are cis-acting
elements of DNA, usually about from 10 to 300 bp, that act on
a promoter to increase its transcription. Many enhancer
sequences are now known from mammalian genes (globin,
elastase, albumin, a-ketoprotein, and insulin). Typically,
however, one will use an enhancer from a eukaryotic cell
virus. Examples include the SV40 enhancer on the late side of
the replication origin (bp 100-270), the cytomegalovirus
early promoter enhancer, the polyoma enhancer on the late
side of the replication origin, and adenovirus enhancers. The
enhancer may be spliced into the vector at a position 5' or 3' to
the fusion protein coding sequence but is preferably located at
a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast,
fungi, insect, plant, animal, human, or nucleated cells from
other multicellular organisms) will also contain sequences
necessary for the termination of transcription and for stabi-
lizing the mRNA. Such sequences are commonly available
from the 5' and occasionally 3' untranslated regions of
eukaryotic or viral DNAs or cDNAs. These regions contain
nucleotide segments transcribed as polyadenylated fragments
in the untranslated portion of the mRNA encoding the fusion
protein.

Various forms of a fusion protein may be recovered from
culture medium or from host cell lysates. If membrane-
bound, it can be released from the membrane using a suitable
detergent solution (e.g., Triton-X 100) or by enzymatic cleav-
age. Cells employed in expression of a fusion protein can be
disrupted by various physical or chemical means, such as
freeze-thaw cycling, sonication, mechanical disruption, or
cell lysing agents.

Once the heterologous fusion proteins of the present inven-
tion are expressed in the appropriate host cell, the analogs can
be isolated and purified. The following procedures are exem-
plary of suitable purification procedures: fractionation on
carboxymethyl cellulose; gel filtration such as Sephadex
G-75; anion exchange resin such as DEAE or Mono-Q; cation
exchange such as CM or Mono-S; metal chelating columns to

14

bind epitope-tagged forms of the polypeptide; reversed-phase
HPLC; chromatofocusing; silica gel; ethanol precipitation;
and ammonium sulfate precipitation.

Various methods of protein purification may be employed
and such methods are known in the art and described, for
example, in Deutscher, *Methods in Enzymology* 182: 83-9
(1990) and Scopes, *Protein Purification: Principles and
Practice*, Springer-Verlag, NY (1982). The purification
step(s) selected will depend on the nature of the production
process used and the particular fusion protein produced. For
example, fusion proteins comprising an Fc fragment can be
effectively purified using a Protein A or Protein G affinity
matrix. Low or high pH buffers can be used to elute the fusion
protein from the affinity matrix. Mild elution conditions will
aid in preventing irreversible denaturation of the fusion pro-
tein.

The heterologous fusion proteins of the present invention
may be formulated with one or more excipients. The fusion
proteins of the present invention may be combined with a
pharmaceutically acceptable buffer, and the pH adjusted to
provide acceptable stability, and a pH acceptable for admin-
istration such as parenteral administration. Optionally, one or
more pharmaceutically-acceptable anti-microbial agents
may be added. Meta-cresol and phenol are preferred pharma-
ceutically-acceptable microbial agents. One or more pharma-
ceutically-acceptable salts may be added to adjust the ionic
strength or tonicity. One or more excipients may be added to
further adjust the isotonicity of the formulation. Glycerin is
an example of an isotonicity-adjusting excipient. Pharmaceu-
tically acceptable means suitable for administration to a
human or other animal and thus, does not contain toxic ele-
ments or undesirable contaminants and does not interfere
with the activity of the active compounds therein.

The heterologous fusion proteins of the present invention
may be formulated as a solution formulation or as a lyo-
philized powder that can be reconstituted with an appropriate
diluent. A lyophilized dosage form is one in which the fusion
protein is stable, with or without buffering capacity to main-
tain the pH of the solution over the intended in-use shelf-life
of the reconstituted product. It is preferable that the solution
comprising the heterologous fusion proteins discussed herein
before lyophilization be substantially isotonic to enable for-
mation of isotonic solutions after reconstitution.

A pharmaceutically-acceptable salt form of the heterolo-
gous fusion proteins of the present invention are within the
scope of the invention. Acids commonly employed to form
acid addition salts are inorganic acids such as hydrochloric
acid, hydrobromic acid, hydriodic acid, sulfuric acid, phos-
phoric acid, and the like, and organic acids such as p-toluene-
sulfonic acid, methanesulfonic acid, oxalic acid, p-bro-
mophenyl-sulfonic acid, carbonic acid, succinic acid, citric
acid, benzoic acid, acetic acid, and the like. Preferred acid
addition salts are those formed with mineral acids such as
hydrochloric acid and hydrobromic acid.

Base addition salts include those derived from inorganic
bases, such as ammonium or alkali or alkaline earth metal
hydroxides, carbonates, bicarbonates, and the like. Such
bases useful in preparing the salts of this invention thus
include sodium hydroxide, potassium hydroxide, ammonium
hydroxide, potassium carbonate, and the like.

The heterologous fusion proteins of the present invention
have biological activity. Biological activity refers to the abil-
ity of the fusion protein to bind to and activate the GLP-1
receptor in vivo and elicit a response. Responses include, but
are not limited to, secretion of insulin, suppression of gluca-
gon, inhibition of appetite, weight loss, induction of satiety,
inhibition of apoptosis, induction of pancreatic beta cell pro-

US 7,452,966 B2

15

liferation, and differentiation of pancreatic beta cells. A representative number of GLP-1 fusion proteins were tested for in vitro as well as in vivo activity. Examples 1 and 2 provide in vitro activity based on the ability of the fusion protein to interact with and activate the human GLP-1 receptor. In both sets of experiments, HEK293 cells over-expressing the human GLP-1 receptor were used. Activation of the GLP-1 receptor in these cells causes adenylyl cyclase activation which in turn induces expression of a reporter gene driven by a cyclic AMP response element (CRE). Example 1 (table 1) provides data wherein the reporter gene is beta lactamase, and example 2 (table 2) provides data wherein the reporter gene is luciferase. Example 3 provides data generated after administration of one of the heterologous fusion proteins of the present invention to rats. Together the data show that the fusion proteins are able to bind to and activate the GLP-1 receptor and appear more potent in vitro than Val⁸-GLP-1(7-37)OH. In addition, the data generated in rats indicate the fusion proteins are active in vivo and have a longer half-life than native GLP-1.

Administration of the heterogeneous fusion proteins may be via any route known to be effective by the physician of ordinary skill. Peripheral parenteral is one such method. Parenteral administration is commonly understood in the medical literature as the injection of a dosage form into the body by a sterile syringe or some other mechanical device such as an infusion pump. Peripheral parenteral routes can include intravenous, intramuscular, subcutaneous, and intraperitoneal routes of administration.

The heterologous fusion proteins of the present invention may also be amenable to administration by oral, rectal, nasal, or lower respiratory routes, which are non-parenteral routes. Of these non-parenteral routes, the lower respiratory route and the oral route are preferred.

The fusion proteins of the present invention can be used to treat a wide variety of diseases and conditions. The fusion proteins of the present invention primarily exert their biological effects by acting at a receptor referred to as the "GLP-1 receptor." Subjects with diseases and/or conditions that respond favorably to GLP-1 receptor stimulation or to the administration of GLP-1 compounds can therefore be treated with the GLP-1 fusion proteins of the present invention. These subjects are said to "be in need of treatment with GLP-1 compounds" or "in need of GLP-1 receptor stimulation". Included are subjects with non-insulin dependent diabetes, insulin dependent diabetes, stroke (see WO 00/16797), myocardial infarction (see WO 98/08531), obesity (see WO 98/19698), catabolic changes after surgery (see U.S. Pat. No. 6,006,753), functional dyspepsia and irritable bowel syndrome (see WO 99/64060). Also included are subjects requiring prophylactic treatment with a GLP-1 compound, e.g., subjects at risk for developing non-insulin dependent diabetes (see WO 00/07617). Subjects with impaired glucose tolerance or impaired fasting glucose, subjects whose body weight is about 25% above normal body weight for the subject's height and body build, subjects with a partial pancreatectomy, subjects having one or more parents with non-insulin dependent diabetes, subjects who have had gestational diabetes and subjects who have had acute or chronic pancreatitis are at risk for developing non-insulin dependent diabetes.

An effective amount of the GLP-1-Fc fusion proteins described herein is the quantity which results in a desired therapeutic and/or prophylactic effect without causing unacceptable side-effects when administered to a subject in need

16

of GLP-1 receptor stimulation. A "desired therapeutic effect" includes one or more of the following: 1) an amelioration of the symptom(s) associated with the disease or condition; 2) a delay in the onset of symptoms associated with the disease or condition; 3) increased longevity compared with the absence of the treatment; and 4) greater quality of life compared with the absence of the treatment. For example, an "effective amount" of a GLP-1-Fc fusion protein for the treatment of diabetes is the quantity that would result in greater control of blood glucose concentration than in the absence of treatment, thereby resulting in a delay in the onset of diabetic complications such as retinopathy, neuropathy or kidney disease. An "effective amount" of a GLP-1-Fc fusion protein for the prevention of diabetes is the quantity that would delay, compared with the absence of treatment, the onset of elevated blood glucose levels that require treatment with anti-hypoglycaemic drugs such as sulfonyl ureas, thiazolidinediones, insulin and/or bisguanidines.

The dose of fusion protein effective to normalize a patient's blood glucose will depend on a number of factors, among which are included, without limitation, the subject's sex, weight and age, the severity of inability to regulate blood glucose, the route of administration and bioavailability, the pharmacokinetic profile of the fusion protein, the potency, and the formulation. Doses may be in the range of 0.01 to 1 mg/kg body weight, preferably in the range of 0.05 to 0.5 mg/kg body weight.

It is preferable that the fusion proteins of the present invention be administered either once every two weeks or once a week. Depending on the disease being treated, it may be necessary to administer the fusion protein more frequently such as two to three time per week.

The present invention will now be described only by way of non-limiting example with reference to the following Examples.

EXAMPLES

Example 1

In Vitro GLP-1 Receptor Activation Assay

HEK-293 cells expressing the human GLP-1 receptor, using a CRE-BLAM system, are seeded at 20,000 to 40,000 cells/well/100 μ l DMEM medium with 10% FBS into a poly-d-lysine coated 96 well black, clear-bottom plate. The day after seeding, the medium is flicked off and 80 μ l plasma-free DMEM medium is added. On the third day after seeding, 20 μ l of plasma-free DMEM medium with 0.5% BSA containing different concentrations of various GLP-1-Fc heterologous fusion protein is added to each well to generate a dose response curve. Generally, fourteen dilutions containing from 3 nanomolar to 30 nanomolar or heterologous GLP-1 Fc fusion protein are used to generate a dose response curve from which EC₅₀ values can be determined. After 5 hours of incubation with the fusion protein, 20 μ l of β -lactamase substrate (CCF2/AM, PanVera LLC) is added and incubation continued for 1 hour at which time fluorescence is determined on a

US 7,452,966 B2

17

cytofluor. The assay is further described in Zlokarnik, et al. (1998), *Science*, 278:84-88. Various GLP-1-Fc fusion proteins are tested and EC₅₀ values are represented in Table 1. The values are relative to values determined for Val⁸-GLP-1 (7-37)OH which is run as an internal control with every experiment.

TABLE 1

Compound	Activ-ity	Std. Dev.
Val ⁸ -GLP-1: Gly ⁸ -Glu ²² -GLP-1(7-37)-2L-IgG4(S228P, F234A, L235A):	100% 301%	99
Gly ⁸ -Glu ²² -GLP-1(7-37)-1.5L-IgG4(S228P, F234A, L235A):	314%	45
Gly ⁸ -Glu ²² -GLP-1(7-37)-1L-IgG4(S228P, F234A, L235A):	468%	120
Gly ⁸ -Glu ²² -Gly ³⁶ -GLP-1(7-37)-2L-IgG4(S228P, F234A, L235A):	441%	35

Example 2

In Vitro GLP-1 Receptor Activation Assay

HEK-293 cells stably expressing the human GLP-1 receptor, using a CRE-Luciferase system, are seeded at 30,000 cells/well/80 µl low serum DMEM F12 medium into 96 well plates. The day after seeding, 20 µl aliquots of test protein dissolved in 0.5% BSA are mixed and incubated with the cells for 5 hours. Generally 12 dilutions containing from 3 pM to 3 nM are prepared at a 5× concentration for each test protein before addition to the cells to generate a dose response curve from which EC₅₀ values are determined. After incubation, 100 µl of Luciferase reagent is added directly to each plate and mixed gently for 2 minutes. Plates are placed in a Tri-lux luminometer and light output resulting from luciferase expression is calculated. Various GLP-1-Fc fusion proteins are tested and EC₅₀ values are represented in Table 2. The values are relative to values determined for Val⁸-GLP-1(7-37)OH which is run as an internal control with every experiment. Because the fusion proteins tested below are dimers, values are corrected taking into account a 2-fold difference in molarity.

TABLE 2

Compound	Activ-ity	Std. Dev.
Val ⁸ -GLP-1: Gly ⁸ -Glu ²² -GLP-1(7-37)-2L-IgG4(S228P, F234A, L235A):	100% 535%	240
Gly ⁸ -Glu ²² -GLP-1(7-37)-1.5L-IgG4(S228P, F234A, L235A):	595%	43
Gly ⁸ -Glu ²² -GLP-1(7-37)-1L-IgG4(S228P, F234A, L235A):	1119%	128
Gly ⁸ -Glu ²² -Gly ³⁶ -GLP-1(7-37)-2L-IgG4(S228P, F234A, L235A):	398%	62
Gly ⁸ -Glu ²² -Gly ³⁶ -GLP-1(7-37)-1L-IgG4(S228P, F234A, L235A):	417%	140

Example 3

Intravenous Glucose Tolerance Test in Rats

The Fc fusion protein, Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-L-IgG4 (S228P,F234A,L235A), is evaluated in an intravenous glucose tolerance test (IVGTT) in rats. At least four rats are

18

included into each of three groups. Group I receives vehicle (table 3), Group II receives 1.79 mg/kg of Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-L-IgG4 (S228P,F234A,L235A) as a single subcutaneous injection (table 4), and Group III receives 0.179 mg/kg of Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-L-IgG4 (S228P, F234A,L235A) as a single subcutaneous injection (table 5). Rats are subcutaneously injected the morning of Day 1. Twenty-four hours following the first injection, 1 µL of glucose (D50) per gram rat body weight is infused as a bolus. Blood samples are taken at 2, 4, 6, 10, 20, and 30 minutes following the bolus infusion of glucose.

TABLE 3

Insulin AUC (ng*min/mL)							
Vehicle:	Rat 1	Rat 2	Rat 3	Rat 4	Rat 5	Average	SEM
0-2	11	9.4	7	11	9.6		
2-4	18.1	9.7	5.6	10.6	8.8		
4-6	13.4	7	3.4	9.6	5.9		
6-10	7.9	3.5	2.5	6	2.9		
10-20	3.7	3	2.4	3	2.4		
20-30	2	0	0	0	2.4		
sum	56.1	32.6	20.9	40.2	32	36.4	5.8

TABLE 4

Insulin AUC (ng*min/mL)							
(1.79 mg/kg)	GLP-1-Fc					Average	SEM
	Rat 1	Rat 2	Rat 3	Rat 4	Rat 5		
0-2	12.3	17.4	16	14	13		
2-4	21.9	13.3	13.2	13.9	13.6		
4-6	16.8	6.5	9.8	11.1	11.7		
6-10	7.6	3.8	9.2	5.8	7.4		
10-20	3	0	0	3.2	5.6		
20-30	0	0	0	0	0		
sum	61.6	41	48.2	48	51.3	50	3.4

TABLE 5

Insulin AUC (ng*min/mL)							
(0.179 mg/kg)	GLP-1-Fc					Average	SEM
	Rat 1	Rat 2	Rat 3	Rat 4			
0-2	14.4	29.2	25.4	23.2			
2-4	13.8	26.3	21.2	21.8			
4-6	11.2	19.4	16.4	15.7			
6-10	6.4	10.6	10.5	8			
10-20	3.6	5.8	5.2	5			
20-30	0	0	0	0			
sum	49.4	91.3	78.7	73.7	78.7	8.7	

Example 4

Pharmacokinetic Study Following a Single Subcutaneous Injection to Cynomolgus Monkeys

A study is performed to characterize the pharmacokinetics (PK) of the Fc fusion protein, Gly⁸-Glu²²-Gly^{6'''}-GLP-1(7-37)-L-IgG4 (S228P,F234A,L235A), when administered as a 0.1 mg/kg by subcutaneous (SC) injection to male cynomolgus monkeys. RIA antibody is specific for the middle portion of GLP. ELISA uses an N-terminus specific capture antibody and an Fc specific detection antibody. Resulting plasma con-

US 7,452,966 B2

19

centrations from both the ELISA and the RIA are used to determine the represented pharmacokinetic parameter values. A representation of the resulting PK parameter values is summarized in table 6. Single-dose SC PK from the RIA is associated with a mean C_{max} of 446.7 ng/mL with a corresponding T_{max} of 17.3 hours. The mean elimination half-life is approximately 79.3 hours (3.3 days). The PK from the ELISA is associated with a mean C_{max} of 292.2 ng/mL with a corresponding T_{max} of 16.7 hours. The mean elimination half-life is approximately 51.6 hours (2.2 days).

TABLE 6

Dose (mg/kg)	Animal #	C_{max}^a (ng/mL)	T_{max}^b (h)	$AUC_{0-\infty}^c$ (ng*h/mL)	$t_{1/2}^d$ (h)	CL/F ^e (mL/h/kg)	Vss/F ^f (mL/kg)
RIA							
0.1	96051	461.0	4.0	37770.5	81.0	2.7	309.2
	96071	430.0	24.0	43150.2	74.2	2.3	248.1
	96091	449.0	24.0	62271.1	82.9	1.6	191.9
RIA	Mean	446.7	17.3	47730.6	79.3	2.2	249.8
	SD	15.6	11.5	12876.5	4.5	0.5	58.7
ELISA							
	96051	315.4	2.0	9062.3	55.2	11.0	879.4
	96071	289.4	24.0	16653.0	50.3	6.0	436.0
	96091	271.9	24.0	19907.4	49.3	5.0	357.0
ELISA	Mean	292.2	16.7	15207.6	51.6	7.3	557.5
	SD	21.9	12.7	5565.2	3.2	3.2	281.6

^aMaximum observed plasma concentration.
^bTime of maximum observed plasma concentration.
^cArea under the plasma concentration-time curve measured from 0 to infinity.
^dElimination half-life.
^eTotal body clearance as a function of bioavailability.
^fVolume of distribution as a function of bioavailability.
SD = Standard deviation.

20

Example 5

Assessment of the Potential Formation of Antibodies Following Repeat Subcutaneous Injections

Designated serum samples from cynomolgus monkeys are tested for the formation of antibodies against Gly⁸-Glu²²-Gly³⁶-GLP-1(7-37)-L-IgG4 (S228P,F234A,L235A) using a direct ELISA format. Microtiter plates are coated with Gly⁸-Glu²²-Gly⁶-GLP-1(7-37)-L-IgG4 (S228P,F234A,L235A) at a 0.1 µg/mL concentration. Monkey serum samples are diluted 50, 500, 1000 and 5000 fold into blocking solution, and 0.05 mL sample/well are incubated approximately one hour. Secondary antibody, Goat<Human Fab'2>-Peroxidase (with 75% cross reactivity to human), is diluted 10,000 fold into block and added at 0.05 ml/well and incubated approximately one hour. Color development using tetramethylbenzidine (TMB) substrate is read at an optical density of 450 nm-630 nm. Duplicate readings are averaged. A GLP-1 antibody was used as a positive control and goat<rabbit>(H+L)-Peroxidase conjugate is the secondary used for detection. Point serum samples are collected prior to dosing, at 24 hours following the second dose, and 168 hours following the first and second SC dose for an evaluation of potential immunogenicity. The presence of antibody titers to G8E22-CEX-L-hIgG4 is interpreted by comparison to predose serum samples and positive control. A representation of the results is presented in table 7.

TABLE 7

Dose 1 Animal#									
Sample Time:	Positive Control	IO7774 Predose	IO7774 168 h	IO7777 Predose	IO7777 168 h	IO7779 Predose	IO7779 168 h	IO7780 Predose	IO7780 168 h
50x	2.854	0.268	0.268	0.160	0.128	0.144	0.152	0.264	0.224
500x	2.270	0.117	0.133	0.052	0.069	0.065	0.061	0.067	0.061
1000x	1.610	0.091	0.075	0.034	0.051	0.047	0.045	0.138	0.049
5000x	0.525	0.056	0.048	0.032	0.037	0.029	0.033	0.051	0.039
Dose 2 Animal#									
Sample Time:	Positive Control	IO7774 Predose	IO7774 24 h	IO7777 Predose	IO7777 24 h	IO7779 Predose	IO7779 24 h	IO7780 Predose	IO7780 24 h
50x	3.056	0.298	0.231	0.164	0.159	0.227	0.176	0.211	0.192
500x	2.247	0.120	0.119	0.048	0.045	0.061	0.060	0.056	0.057
1000x	1.673	0.090	0.086	0.039	0.041	0.046	0.045	0.043	0.048
5000x	0.534	0.039	0.042	0.030	0.034	0.033	0.036	0.033	0.034
Dose 1 Animal#									
Sample Time:	Positive Control	IO7774 Predose	IO7774 168 h	IO7777 Predose	IO7777 168 h	IO7779 Predose	IO7779 168 h	IO7780 Predose	IO7780 168 h
50x	3.075	0.413	0.270	0.174	0.182	0.185	0.190	0.224	0.191
500x	2.173	0.097	0.103	0.042	0.051	0.056	0.057	0.048	0.053
1000x	1.510	0.066	0.067	0.038	0.040	0.037	0.046	0.043	0.043
5000x	0.474	0.042	0.042	0.033	0.046	0.033	0.033	0.036	0.041

Pharmacodynamic Study Following a Single Subcutaneously Injection to Cynomolgus Monkeys in the Fasting State and During a Graded Intravenous Glucose Infusion

In Phase 1 (Study Day 1) a subcutaneous injection of vehicle is administered. A graded intravenous glucose (20% dextrose) infusion of 5, 10, and 25 mg/kg/min is then administered immediately after the vehicle injection. In Phase 2 (Study Day 3), a subcutaneous injection of a GLP-1 fusion protein (0.1 mg/kg) is administered. In Phase 3, a graded intravenous glucose infusion is performed approximately 96 hours following the GLP-1 fusion injection.

Graded intravenous glucose infusion procedures are conducted in sedated monkeys after a 16-hr overnight fast. For both intravenous glucose infusions, baseline samples will be drawn every 10 min for 20 min to define baseline. A stepped-up glucose infusion is initiated at +20 min at a rate of 5 mg/kg/min, followed by infusions of 10 mg/kg/min, and 25 mg/kg/min. Each infusion rate is administered for a period of 20 minutes. Blood samples are taken at 10 minute intervals for measurement of glucose, insulin, and glucagon. Approximately 1.0 mL of blood is collected at -20, -10 min, 0 pre-glucose infusions, and at 10, 20, 30, 40, 50, and 60 minutes following glucose infusion for Phases 1 and 3.

A representation of the data are shown in table 8.

TABLE 8

Glucose AUC					
AUC			AUC		
Group	Animal	(min*mg/dL)	Group	Animal	(min*mg/dL)
GLP-Fc	9423	7447	vehicle	9423	8077
	9424	7470		9424	15006
	9510	5153		9510	7116
	9513	6303		9513	7459
	9516	5413		9516	8728
	9530	5240		9530	7863
					N 6
Mean		6171	Mean		9041

TABLE 8-continued

	SD	1078		SD	2973
	SE	440		SE	1214
<u>Insulin AUC</u>					
		AUC			AUC
Group	Animal	(min*ng/mL)	Group	Animal	(min*ng/mL)
GLP-Fc	9423	129	vehicle	9423	38
	9424	138		9424	29
	9510	357		9510	69
	9513	161		9513	64
	9516	376		9516	38
	9530	215		9530	68
	Mean	229		Mean	51
	SD	111		SD	18
	SE	45		SE	7

Glucagon levels were not statistically different between the vehicle and the GLP-1 fusion protein dosed monkeys

Example 7

Pharmacodynamic Study Following Single Subcutaneously Injections of Three Different Doses to Rats in the Fasting State and During a Graded Intravenous Glucose Infusion

Chronically cannulated rats are assigned to either vehicle control (saline) or one of 3 treatment groups (GLP-1 fusion protein; 0.0179 mg/kg, 0.179 mg/kg, or 1.79 mg/kg). The GLP-1 fusion protein and vehicle are administered via subcutaneous injection. Twenty-four hours after treatment, overnight fasted (16 h) rats are subjected to a graded intravenous glucose infusion test. The graded glucose infusion test consists of a baseline saline infusion period (20 min), followed by two 30 min glucose infusion phases at 5 and 15 mg/kg/min, respectively. Plasma samples are collected at -20, -10 min, 0 pre-glucose infusions (baseline), and at 10, 20, 30, 40, 50, and 60 minutes.

A representation of the data are shown in table 9.

TABLE 9

5 mg/Kg/min			15 mg/Kg/min		
Vehicle	4.3 ± 0.2 (n = 18)		12.7 ± 0.9 (n = 18)		
0.0179 mg/Kg	5.6 ± 0.4 (n = 4)		15.9 ± 1.8 (n = 4)		
0.179 mg/Kg	9.0 ± 1.1* (n = 6)		28.0 ± 3.8* (n = 6)		
1.79 mg/Kg	20.5 ± 3.0* (n = 4)		52.7 ± 7.2* (n = 4)		

*P ≤ 0.05 versus vehicle

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 8

<210> SEQ ID NO 1
<211> LENGTH: 230
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)

-continued

```
<223> OTHER INFORMATION: Xaa at position 1 is Ala or absent
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (16)..(16)
<223> OTHER INFORMATION: Xaa at position 16 is Pro or Glu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (17)..(17)
<223> OTHER INFORMATION: Xaa at position 17 is Phe, Val, or Ala
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: Xaa at position 18 is Leu, Glu, or Ala
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (80)..(80)
<223> OTHER INFORMATION: Xaa at position 80 is Asn or Ala
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (230)..(230)
<223> OTHER INFORMATION: Xaa at position 230 is Lys or is absent

<400> SEQUENCE: 1

Xaa Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Xaa
1          5          10          15

Xaa Xaa Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
          20          25          30

Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
          35          40          45

Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly
          50          55          60

Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Xaa
65          70          75          80

Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp
          85          90          95

Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro
          100          105          110

Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu
          115          120          125

Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn
          130          135          140

Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
145          150          155          160

Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
          165          170          175

Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg
          180          185          190

Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys
          195          200          205

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
          210          215          220

Ser Leu Ser Leu Gly Xaa
225          230

<210> SEQ ID NO 2
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 2
```


-continued

Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser									
1				5					10					15									
<210> SEQ ID NO 3																							
<211> LENGTH: 6																							
<212> TYPE: PRT																							
<213> ORGANISM: Homo sapiens																							
<400> SEQUENCE: 3																							
Pro	Pro	Cys	Pro	Ser	Cys																		
1				5																			
<210> SEQ ID NO 4																							
<211> LENGTH: 22																							
<212> TYPE: PRT																							
<213> ORGANISM: Artificial																							
<220> FEATURE:																							
<223> OTHER INFORMATION: Synthetic Construct																							
<400> SEQUENCE: 4																							
Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly									
1				5					10					15									
Ser	Gly	Gly	Gly	Gly	Ser																		
				20																			
<210> SEQ ID NO 5																							
<211> LENGTH: 825																							
<212> TYPE: DNA																							
<213> ORGANISM: Homo sapiens																							
<400> SEQUENCE: 5																							
cacggcgagg gcaccttcac ctccgacgtg tcctcctatc tcgaggagca ggccgccaag													60										
gaattcatcg cctggctggt gaagggcggc ggcggtggtg gtggctccgg aggcggcggc													120										
tctggtggcg gtggcagcgc tgagtccaaa tatgggtcccc catgcccacc ctgcccagca													180										
cctgaggccg ccggggggacc atcagtcttc ctgttcccc caaaacccaa ggacactctc													240										
atgatctccc ggacccctga ggtcacgtgc gtggtggtgg acgtgagcca ggaagacccc													300										
gaggtccagt tcaactggta cgtggatggc gtggaggtgc ataatgcaa gacaaagccg													360										
cgggaggagc agttcaacag cacgtaccgt gtggtcagcg tcctcaccgt cctgcaccag													420										
gactggctga acggcaagga gtacaagtgc aaggtctcca acaaaggcct cccgtcctcc													480										
atcgagaaaa ccatctccaa agccaaaggg cagccccgag agccacaggt gtacaccctg													540										
cccccatccc aggaggagat gaccaagaac caggtcagcc tgacctgcct ggtcaaaggc													600										
ttctacccca gcgacatcgc cgtggagtgg gaaagcaatg ggcagccgga gaacaactac													660										
aagaccacgc ctcccgtgct ggactccgac ggctccttct tcctctacag caggctaacc													720										
gtggacaaga gcaggtggca ggaggggaat gtcttctcat gctccgtgat gcatgaggct													780										
ctgcacaacc actacacaca gaagagcctc tccctgtctc tgggt													825										
<210> SEQ ID NO 6																							
<211> LENGTH: 30																							
<212> TYPE: PRT																							
<213> ORGANISM: Artificial																							
<220> FEATURE:																							
<223> OTHER INFORMATION: Synthetic Construct																							
<400> SEQUENCE: 6																							
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser									
1				5					10					15									

- continued

Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser		
			20					25					30		
<210> SEQ ID NO 7															
<211> LENGTH: 25															
<212> TYPE: PRT															
<213> ORGANISM: Artificial															
<220> FEATURE:															
<223> OTHER INFORMATION: Synthetic Construct															
<400> SEQUENCE: 7															
Asp	Ala	Ala	Ala	Lys	Glu	Ala	Ala	Ala	Lys	Asp	Ala	Ala	Ala	Arg	Glu
1				5					10					15	
Ala	Ala	Ala	Arg	Asp	Ala	Ala	Ala	Lys							
			20					25							
<210> SEQ ID NO 8															
<211> LENGTH: 14															
<212> TYPE: PRT															
<213> ORGANISM: Artificial															
<220> FEATURE:															
<223> OTHER INFORMATION: Synthetic Construct															
<400> SEQUENCE: 8															
Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg		
1				5				10							

We claim:	30	and a peptide linker comprising SEQ ID NO:8
1. A heterologous fusion protein comprising:		Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-
a GLP-1 analog comprising SEQ ID NO:1		Gly-Gly-Ser
His-Xaa ₈ -Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-		wherein the N-terminal glycine of the peptide linker is
Tyr-Leu-Glu-Glu-Gln-Ala-Ala-Lys-Glu-Phe-Ile-	35	directly fused to the C-terminal glycine residue of the
Ala-Trp-Leu-Val-Lys-Gly-Gly-Gly		GLP-1 analog and the C-terminal serine of the peptide
wherein Xaa ₈ is Gly;		linker is directly fused to the N-terminal alanine of the
an Fc portion of an immunoglobulin comprising SEQ ID		Fc portion.
NO:7		2. A heterologous fusion protein comprising:
Ala-Glu-Ser-Lys-Tyr-Gly-Pro-Pro-Cys-Pro-Pro-Cys-	40	a GLP-1 analog comprising SEQ ID NO:1
Pro-Ala-Pro-Xaa ₁₆ -Xaa ₁₇ -Xaa ₁₈ -Gly-Gly-Pro-Ser-		His-Xaa ₈ -Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-
Val-Phe-Leu-Phe-Pro-Pro-Lys-Pro-Lys-Asp-Thr-		Tyr-Leu-Glu-Glu-Gln-Ala-Ala-Lys-Glu-Phe-Ile-
Leu-Met-Ile-Ser-Arg-Thr-Pro-Glu-Val-Thr-Cys-		Ala-Trp-Leu-Val-Lys-Gly-Gly-Gly
Val-Val-Val-Asp-Val-Ser-Gln-Glu-Asp-Pro-Glu-		wherein Xaa ₈ is Gly;
Val-Gln-Phe-Asn-Trp-Tyr-Val-Asp-Gly-Val-Glu-	45	an Fc portion of an immunoglobulin comprising SEQ ID
Val-His-Asn-Ala-Lys-Thr-Lys-Pro-Arg-Glu-Glu-		NO:7
Gln-Phe-Xaa ₈₀ -Ser-Thr-Tyr-Arg-Val-Val-Ser-Val-		Ala-Glu-Ser-Lys-Tyr-Gly-Pro-Pro-Cys-Pro-Pro-Cys-
Leu-Thr-Val-Leu-His-Gln-Asp-Trp-Leu-Asn-Gly-		Pro-Ala-Pro-Xaa ₁₆ -Xaa ₁₇ -Xaa ₁₈ -Gly-Gly-Pro-Ser-
Lys-Glu-Tyr-Lys-Cys-Lys-Val-Ser-Asn-Lys-Gly-		Val-Phe-Leu-Phe-Pro-Pro-Lys-Pro-Lys-Asp-Thr-
Leu-Pro-Ser-Ser-Ile-Glu-Lys-Thr-Ile-Ser-Lys-Ala-	50	Leu-Met-Ile-Ser-Arg-Thr-Pro-Glu-Val-Thr-Cys-
Lys-Gly-Gln-Pro-Arg-Glu-Pro-Gln-Val-Tyr-Thr-		Val-Val-Val-Asp-Val-Ser-Gln-Glu-Asp-Pro-Glu-
Leu-Pro-Pro-Ser-Gln-Glu-Glu-Met-Thr-Lys-Asn-		Val-Gln-Phe-Asn-Trp-Tyr-Val-Asp-Gly-Val-Glu-
Gln-Val-Ser-Leu-Thr-Cys-Leu-Val-Lys-Gly-Phe-		Val-His-Asn-Ala-Lys-Thr-Lys-Pro-Arg-Glu-Glu-
Tyr-Pro-Ser-Asp-Ile-Ala-Val-Glu-Trp-Glu-Ser-Asn-		Gln-Phe-Xaa ₈₀ -Ser-Thr-Tyr-Arg-Val-Val-Ser-Val-
Gly-Gln-Pro-Glu-Asn-Asn-Tyr-Lys-Thr-Thr-Pro-	55	Leu-Thr-Val-Leu-His-Gln-Asp-Trp-Leu-Asn-Gly-
Pro-Val-Leu-Asp-Ser-Asp-Gly-Ser-Phe-Phe-Leu-		Lys-Glu-Tyr-Lys-Cys-Lys-Val-Ser-Asn-Lys-Gly-
Tyr-Ser-Arg-Leu-Thr-Val-Asp-Lys-Ser-Arg-Trp-		Leu-Pro-Ser-Ser-Ile-Glu-Lys-Thr-Ile-Ser-Lys-Ala-
Gln-Glu-Gly-Asn-Val-Phe-Ser-Cys-Ser-Val-Met-		Lys-Gly-Gln-Pro-Arg-Glu-Pro-Gln-Val-Tyr-Thr-
His-Glu-Ala-Leu-His-Asn-His-Tyr-Thr-Gln-Lys-		Leu-Pro-Pro-Ser-Gln-Glu-Glu-Met-Thr-Lys-Asn-
Ser-Leu-Ser-Leu-Ser-Leu-Gly-Xaa ₂₃₀	60	Gln-Val-Ser-Leu-Thr-Cys-Leu-Val-Lys-Gly-Phe-
wherein:		Tyr-Pro-Ser-Asp-Ile-Ala-Val-Glu-Trp-Glu-Ser-Asn-
Xaa at position 16 is Glu;		Gly-Gln-Pro-Glu-Asn-Asn-Tyr-Lys-Thr-Thr-Pro-
Xaa at position 17 is Ala;		Pro-Val-Leu-Asp-Ser-Asp-Gly-Ser-Phe-Phe-Leu-
Xaa at position 18 is Ala;		Tyr-Ser-Arg-Leu-Thr-Val-Asp-Lys-Ser-Arg-Trp-
Xaa at position 80 is Asn; and	65	Gln-Glu-Gly-Asn-Val-Phe-Ser-Cys-Ser-Val-Met-
Xaa at position 230 is Lys;		His-Glu-Ala-Leu-His-Asn-His-Tyr-Thr-Gln-Lys-
		Ser-Leu-Ser-Leu-Ser-Leu-Gly-Xaa ₂₃₀

US 7,452,966 B2

29

wherein:
Xaa at position 16 is Glu;
Xaa at position 17 is Ala;
Xaa at position 18 is Ala;
Xaa at position 80 is Asn; and
Xaa at position 230 is absent;
and a peptide linker comprising SEQ ID NO:8
Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-
Gly-Gly-Ser
wherein the N-terminal glycine of the peptide linker is
directly fused to the C-terminal glycine residue of the
GLP-1 analog and the C-terminal serine of the peptide
linker is directly fused to the N-terminal alanine of the
Fc portion.
3. A heterologous fusion protein whose amino acid
sequence consists of:
a GLP-1 analog consisting of SEQ ID NO:1
His-Xaa₈-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-
Tyr-Leu-Glu-Glu-Gln-Ala-Ala-Lys-Glu-Phe-Ile-
Ala-Trp-Leu-Val-Lys-Gly-Gly-Gly
wherein Xaa₈ is Gly;
an Fc portion of an immunoglobulin consisting of SEQ ID
NO:7
Ala-Glu-Ser-Lys-Tyr-Gly-Pro-Pro-Cys-Pro-Pro-Cys-
Pro-Ala-Pro-Xaa₁₆-Xaa₁₇-Xaa₁₈-Gly-Gly-Pro-Ser-
Val-Phe-Leu-Phe-Pro-Pro-Lys-Pro-Lys-Asp-Thr-
Leu-Met-Ile-Ser-Arg-Thr-Pro-Glu-Val-Thr-Cys-
Val-Val-Val-Asp-Val-Ser-Gln-Glu-Asp-Pro-Glu-
Val-Gln-Phe-Asn-Trp-Tyr-Val-Asp-Gly-Val-Glu-
Val-His-Asn-Ala-Lys-Thr-Lys-Pro-Arg-Glu-Glu-
Gln-Phe-Xaa₈₀-Ser-Thr-Tyr-Arg-Val-Val-Ser-Val-
Leu-Thr-Val-Leu-His-Gln-Asp-Trp-Leu-Asn-Gly-
Lys-Glu-Tyr-Lys-Cys-Lys-Val-Ser-Asn-Lys-Glu-
Leu-Pro-Ser-Ser-Ile-Glu-Lys-Thr-Ile-Ser-Lys-Ala-
Lys-Gly-Gln-Pro-Arg-Glu-Pro-Gln-Val-Tyr-Thr-
Leu-Pro-Pro-Ser-Gln-Glu-Glu-Met-Thr-Lys-Asn-
Gln-Val-Ser-Leu-Thr-Cys-Leu-Val-Lys-Gly-Phe-
Tyr-Pro-Ser-Asp-Ile-Ala-Val-Glu-Trp-Glu-Ser-Asn-
Gly-Gln-Pro-Glu-Asn-Asn-Tyr-Lys-Thr-Thr-Pro-
Pro-Val-Leu-Asp-Ser-Asp-Gly-Ser-Phe-Phe-Leu-
Tyr-Ser-Arg-Leu-Thr-Val-Asp-Lys-Ser-Arg-Trp-
Gln-Glu-Gly-Asn-Val-Phe-Ser-Cys-Ser-Val-Met-
His-Glu-Ala-Leu-His-Asn-His-Tyr-Thr-Gln-Lys-
Ser-Leu-Ser-Leu-Ser-Leu-Gly-Xaa₂₃₀
wherein:
Xaa at position 16 is Glu;
Xaa at position 17 is Ala;
Xaa at position 18 is Ala;
Xaa at position 80 is Asn; and
Xaa at position 230 is Lys;
and a peptide linker consisting of SEQ ID NO:8
Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-
Gly-Gly-Ser

30

wherein the N-terminal glycine of the peptide linker is
directly fused to the C-terminal glycine residue of the
GLP-1 analog and the C-terminal serine of the peptide
linker is directly fused to the N-terminal alanine of the
Fc portion.
4. A heterologous fusion protein consisting of:
a GLP-1 analog consisting of SEQ ID NO:1
His-Xaa₈-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-
Tyr-Leu-Glu-Glu-Gln-Ala-Ala-Lys-Glu-Phe-Ile-
Ala-Trp-Leu-Val-Lys-Gly-Gly-Gly
wherein Xaa₈ is Gly;
an Fc portion of an immunoglobulin consisting of SEQ ID
NO:7
Ala-Glu-Ser-Lys-Tyr-Gly-Pro-Pro-Cys-Pro-Pro-Cys-
Pro-Ala-Pro-Xaa₁₆-Xaa₁₇-Xaa₁₈-Gly-Gly-Pro-Ser-
Val-Phe-Leu-Phe-Pro-Pro-Lys-Pro-Lys-Asp-Thr-
Leu-Met-Ile-Ser-Arg-Thr-Pro-Glu-Val-Thr-Cys-
Val-Val-Val-Asp-Val-Ser-Gln-Glu-Asp-Pro-Glu-
Val-Gln-Phe-Asn-Trp-Tyr-Val-Asp-Gly-Val-Glu-
Val-His-Asn-Ala-Lys-Thr-Lys-Pro-Arg-Glu-Glu-
Gln-Phe-Xaa₈₀-Ser-Thr-Tyr-Arg-Val-Val-Ser-Val-
Leu-Thr-Val-Leu-His-Gln-Asp-Trp-Leu-Asn-Gly-
Lys-Glu-Tyr-Lys-Cys-Lys-Val-Ser-Asn-Lys-Gly-
Leu-Pro-Ser-Ser-Ile-Glu-Lys-Thr-Ile-Ser-Lys-Ala-
Lys-Gly-Gln-Pro-Arg-Glu-Pro-Gln-Val-Tyr-Thr-
Leu-Pro-Pro-Ser-Gln-Glu-Glu-Met-Thr-Lys-Asn-
Gln-Val-Ser-Leu-Thr-Cys-Leu-Val-Lys-Gly-Phe-
Tyr-Pro-Ser-Asp-Ile-Ala-Val-Glu-Trp-Glu-Ser-Asn-
Gly-Gln-Pro-Glu-Asn-Asn-Tyr-Lys-Thr-Thr-Pro-
Pro-Val-Leu-Asp-Ser-Asp-Gly-Ser-Phe-Phe-Leu-
Tyr-Ser-Arg-Leu-Thr-Val-Asp-Lys-Ser-Arg-Trp-
Gln-Glu-Gly-Asn-Val-Phe-Ser-Cys-Ser-Val-Met-
His-Glu-Ala-Leu-His-Asn-His-Tyr-Thr-Gln-Lys-
Ser-Leu-Ser-Leu-Ser-Leu-Gly-Xaa₂₃₀
wherein:
Xaa at position 16 is Glu;
Xaa at position 17 is Ala;
Xaa at position 18 is Ala;
Xaa at position 80 is Asn; and
Xaa at position 230 is absent;
and a peptide linker consisting of SEQ ID NO:8
Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser-Gly-Gly-
Gly-Gly-Ser
wherein the N-terminal glycine of the peptide linker is
directly fused to the C-terminal glycine residue of the
GLP-1 analog and the C-terminal serine of the peptide
linker is directly fused to the N-terminal alanine of the
Fc portion.
5. The heterologous fusion protein of claim 4 wherein the
fusion protein is encoded by the DNA of SEQ ID NO:20.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE

(12) CERTIFICATE EXTENDING PATENT TERM
UNDER 35 U.S.C. § 156

(68) PATENT NO. : 7,452,966
(45) ISSUED : November 18, 2008
(75) INVENTOR : Glaesner et al.
(73) PATENT OWNER : Eli Lilly and Company
(95) PRODUCT : TRULICITY® (dulaglutide)

This is to certify that an application under 35 U.S.C. § 156 has been filed in the United States Patent and Trademark Office, requesting extension of the term of U.S. Patent No. 7,452,966 based upon the regulatory review of the product TRULICITY® (dulaglutide) by the Food and Drug Administration. According to United States Patent and Trademark Office records, the original expiration date of the patent as of the date of issuance of this certificate is July 06, 2024. Because it appears that the requirements of the law have been met, this certificate extends the term of the patent for the period of

(94) 1,249 days

subject to the payment of maintenance fees as provided by law, with all rights pertaining thereto as provided by 35 U.S.C. § 156.

I have caused the seal of the United States Patent and Trademark Office to be affixed this 15th day of August 2018.



Andrei Iancu

Andrei Iancu
Under Secretary of Commerce for Intellectual Property and
Director of the United States Patent and Trademark Office



US007452966C1

(12) **EX PARTE REEXAMINATION CERTIFICATE** (8835th)
United States Patent
Glaesner et al.

(10) **Number:** **US 7,452,966 C1**
(45) **Certificate Issued:** ***Jan. 31, 2012**

(54) **GLP-1 ANALOG FUSION PROTEINS**

(75) Inventors: **Wolfgang Glaesner**, Indianapolis, IN (US); **Rohn Lee Millican, Jr.**, Indianapolis, IN (US); **Andrew Mark Vick**, Fishers, IN (US)

(73) Assignee: **Eli Lilly and Company**, Indianapolis, IN (US)

Reexamination Request:

No. 90/011,778, Jun. 30, 2011

Reexamination Certificate for:

Patent No.: **7,452,966**
Issued: **Nov. 18, 2008**
Appl. No.: **10/558,627**
Filed: **Nov. 29, 2005**

(*) Notice: This patent is subject to a terminal disclaimer.

(22) PCT Filed: **Jun. 10, 2004**

(86) PCT No.: **PCT/US2004/015595**

§ 371 (c)(1),
(2), (4) Date: **Nov. 29, 2005**

(87) PCT Pub. No.: **WO2005/000892**

PCT Pub. Date: **Jan. 6, 2005**

Related U.S. Application Data

(60) Provisional application No. 60/477,880, filed on Jun. 12, 2003.

(51) **Int. Cl.**

A61K 38/00 (2006.01)

A61K 38/28 (2006.01)

A61P 3/10 (2006.01)

C07K 14/50 (2006.01)

C07K 14/605 (2006.01)

(52) **U.S. Cl.** **530/324; 530/308**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

To view the complete listing of prior art documents cited during the proceeding for Reexamination Control Number 90/011,778, please refer to the USPTO's public Patent Application Information Retrieval (PAIR) system under the Display References tab.

Primary Examiner—Gary Kunz

(57) **ABSTRACT**

The invention provides specific GLP-1 analogs fused to specific IgG4-Fc derivatives. These fusion proteins have an increased half-life, decreased immunogenicity, and reduce effector activity. The fusion proteins are useful in treating diabetes, obesity, irritable bowel syndrome and other conditions that would be benefited by lowering plasma glucose, inhibiting gastric and/or intestinal motility and inhibiting gastric and/or intestinal emptying, or inhibiting food intake.

US 7,452,966 C1

1
EX PARTE
REEXAMINATION CERTIFICATE
ISSUED UNDER 35 U.S.C. 307

NO AMENDMENTS HAVE BEEN MADE TO
THE PATENT

2
AS A RESULT OF REEXAMINATION, IT HAS BEEN
DETERMINED THAT:

5 The patentability of claims **1-5** is confirmed.

* * * * *